

Chesapeake Bay Hypoxic Volume Forecasts

Donald Scavia¹, Isabella Bertani²

¹University of Michigan


²Chesapeake Bay Program

June 23, 2021

In 2020 we transitioned research forecasts developed at the University of Michigan to operational forecasts by the Chesapeake Bay Program. We are pleased to move our research model into operations.

The 2021 Forecast - Given the average January-May 2021 total nitrogen load of 314,693 kg/day, the forecasts are 7.9 km³ (95% prediction interval: 3.7-13.3) for the July average, 5.2 km³ (3.1-7.2) for the summer average, and 690 km³-days (430-940) for the total annual hypoxia. All three are close to their long-term averages.

End-of-summer update: hypoxic volumes estimated from monitoring data were 5.5 km³ (July average), 5.4 km³ (summer average) and 694 km³ (total annual).



Forecasting relationship of total nitrogen vs. hypoxic volume (left: total annual HV, center: summer average HV, right: July average HV). Solid black curves are response curves calibrated to three sets of observations over 1985-2018 (means of the three sets of observations shown as circles). Shaded gray areas are 95% confidence intervals and solid gray curves are 95% prediction intervals. The black vertical line represents the 2021 Jan-May TN load of 314,693 kg/day and the black horizontal lines show the forecast hypoxic volume and prediction intervals associated with this load. The red horizontal dashed lines indicate the long-term average total annual and summer average hypoxic volume predicted by the model if load reductions established by the WIP3 Planning Targets (WIP 2020) and consistent with the Bay's TMDL (US EPA 2010) are achieved.


Model Update and Transition – Up to 2019, the model relied on Susquehanna River total nitrogen (TN) load as a proxy for the Bay's entire load to predict average July hypoxic volume (HV). Updates between 2007 and 2019 were based primarily on model recalibrations. The 2007 and 2008 model was calibrated to 1950-2003 data from Hagy (2002), and after that to HV from the Chesapeake Bay Program (CBP). The 2010-2014 model was calibrated to rolling 3-year



windows (Evans and Scavia 2011), and the 2015–2019 models were calibrated to the full 1985–2017 data set and updated nitrogen loads from the USGS.

During the 2020 transition, the model was calibrated to a broader set of more relevant loads and HV estimates (Scavia et al. 2021). Using loads from all of the primary tributaries, as well as point sources downstream from the tributary monitoring sites, the new forecasts include summer average and total annual HV, in addition to the July average. Summer average HV reduces the impact of individual storms that can disrupt individual cruise data, and total annual HV is a strong integrator of the full impact of the load.

Forecast Track Record – Annual forecasts provided each spring since 2007 have been shown to be rather robust (Testa et al. 2017). Between 2007 and 2018, observed HV fell within the forecast credible intervals in 8 of the 12 years with the under-prediction in 2011 and over-prediction in 2008 just outside the intervals. The substantial over-predictions in 2007 and 2014 were likely because storms like Hurricane Arthur disrupted stratification and temporarily reduced the observed July HV. To circumvent this, we included summer average and total annual HV beginning with the 2020 forecast.


To establish a comparable track record for the new model, a series of blind forecasts were made for the same time period. For each year, the model was calibrated using data up to the preceding year and then used to forecast that year’s HV.





Observed total annual (top), summer average (center) and july average (bottom) hypoxic volume for the period of record. Green bars represent the observed HV (+/- 95% confidence interval), yellow represent blind forecasts (+/- 95% prediction error), and orange represent actual forecasts generated with the most updated model version. The red bar is this year's forecast and the black horizontal line is the long-term average. The red horizontal dashed lines indicate the long-term average total annual and summer average hypoxic volume predicted by the model if load reductions established by the WIP3 Planning Targets (WIP 2020) and consistent with the Bay's TMDL (US EPA 2010) are achieved.

Nitrogen Loads – The nitrogen load, one of the key drivers of hypoxia in the Bay, increased significantly since the 1950s and remains relatively constant but highly variable from year to year. The plot below shows the Jan-May TN loads from primary tributaries and downstream point sources used to drive the model.



The Jan-May TN load (kg/day) including point sources downstream from the tributary monitoring stations (PS), and combined loads from the Susquehanna, Rappahannock, Potomac, Patuxent, Pamunkey, Mattaponi, Appomattox, James and Choptank (RIM). The horizontal red dashed line indicates the long-term average Jan-May TN load.

The Model - The model is an adaptation of the Streeter-Phelps model that simulates dissolved oxygen (DO) depletion in rivers downstream from a point source of organic matter (Streeter and Phelps 1925). It has been applied extensively to rivers and estuaries (Chapra 1997), as well as to the northern Gulf of Mexico (Scavia et al. 2003, 2004, 2006, 2017, 2019) and the Chesapeake Bay (Scavia et al. 2006, Liu et al. 2011, Evans and Scavia 2010).

$$DO = DO_s - \frac{k_d BOD_u F}{k_r - k_d} (e^{-k_d \frac{x}{v}} - e^{-k_r \frac{x}{v}}) - D_i e^{-k_r \frac{x}{v}}$$

where DO = dissolved oxygen (mg/l), DO_s = oxygen saturation (mg/l), k_d = organic matter decay coefficient (1/day), k_r = reaeration coefficient (1/day), BOD_u = initial organic matter (mg/l), x = upstream distance (km), F = fraction of organic matter sinking below the pycnocline (unitless), D_i = initial oxygen deficit (mg l), and v = net advection (km/day). While in the original Streeter-Phelps formulation v represents river advection, here it is a parameterization of the combined effects of horizontal transport and all ecological processes resulting in subsequent settling of organic matter from the surface. Therefore, it is a bulk parameter with no simple physical analog.

Nitrogen load is used as a surrogate for organic matter deposited below the pycnocline at the model origin (220 km down bay from the Susquehanna River mouth) with model distance following the up-estuary flow of bottom water. Simulated HV in year i (V_i) is calculated from

the overall length (L_i) of the DO profile with concentrations less than 2 mg/l using the following empirical relationship derived from Chesapeake Bay measurements (Scavia et al. 2006):

$$V_i = 0.00391 * L_i^2$$

Other assumptions of the model include: transport results from advection rather than longitudinal dispersion, subpycnocline oxygen consumption can be modeled as a first-order process proportional to organic matter concentration, oxygen flux across the pycnocline can be modeled as a first-order process proportional to the difference between surface and bottom layer oxygen concentrations, and subpycnocline organic matter oxygen demand is proportional to TN load. Tests of these assumptions, as well as calibration to mid-July subpycnocline oxygen concentration profiles and HVs from 1950 to 2003, have been described elsewhere (Scavia et al. 2006).

Bayesian Inference – The original model (Scavia et al. 2006) was a Monte Carlo implementation used to accommodate potential variation in the bulk parameter, v . It was subsequently reformulated within a Bayesian framework (Liu et al. 2011, Evans and Scavia 2010) to account for uncertainty in additional parameters. In the present version, the model was calibrated using Bayesian fitting conducted with the software WinBUGS version 1.4.3 (Lunn et al. 2000, Gill 2002, Gelman and Hill 2007) interfaced with R version 3.5.2 (R Core Team 2018) through the package R2WinBUGS version 2.1-21 (Sturtz et al. 2005). All model parameters were kept constant across years and given the same priors used in the most recent model applications (Evans and Scavia 2010; Liu et al. 2011). We ran four Markov Chain Monte Carlo chains with 5,000 iterations each and checked convergence by ensuring that $\hat{R} < 1.1$ for all model parameters. We assessed model performance using the coefficient of determination (R^2), Nash-Sutcliffe Efficiency (NSE), root mean square error (RMSE), mean absolute error (MAE), and residual standard deviation (RSTDE). Details are in Scavia et al. (2021).

References

- Chapra, S.C., 1997. Surface Water-Quality Modeling. McGraw-Hill, New York.
- Evans, M.A. and D. Scavia 2011. Forecasting hypoxia in the Chesapeake Bay and Gulf of Mexico: Model accuracy, precision, and sensitivity to ecosystem change. *Environ. Res. Letters*. doi:10.1088/1748-9326/6/1/015001
- Liu, Y., Arhonditsis, G.B., Stow, C.A., Scavia, D., 2011. Predicting the hypoxic-volume in Chesapeake Bay with the Streeter Phelps model: a Bayesian approach. *J. Am. Water Resour. Assoc.* 1 (6), 1348–1363 DOI: 10.1111/j.1752-1688.2011.00588.x
- Hagy, J. D., 2002. Eutrophication, hypoxia and trophic transfer efficiency in Chesapeake Bay. PhD dissertation, University of Maryland at College Park, College Park, Maryland.
- Qian, S. S., C. A. Stow, and M. E. Borsuk. 2003. On Monte Carlo methods for Bayesian inference. *Ecological Modelling*, 159(2-3): 269-277.
- Scavia, D., I. Bertani, C. Long, and Y. Wang. 2019a. Chesapeake Bay Hypoxic Volume Forecasts. June 7, 2019. Available at: <http://scavia.seas.umich.edu/wp-content/uploads/2019/06/2019-Chesapeake-Bay-forecast.pdf>
- Scavia, D., I. Bertani, D.R. Obenour, R.E. Turner, D.R. Forrest, A. Katin. 2017 Ensemble modeling informs hypoxia management in the northern Gulf of Mexico. *Proc. Nat. Acad. Sci.* 114:8823-8828

- Scavia D., Bertani I., Testa J.M., Bever A.J., Blomquist J.D., Friedrichs M.A.M., Linker L.C., Michael B.D., Murphy R.R., Shenk G.W. 2021. Advancing Estuarine Ecological Forecasts: Seasonal Hypoxia in Chesapeake Bay. *Ecological Applications*.
<https://doi.org/10.1002/eap.2384>
- Scavia, D., D. Justic, D. Obenour, K. Craig, L. Wang. 2019. Hypoxic volume is more responsive than hypoxic area to nutrient load reductions in the northern Gulf of Mexico – and it matters to fish and fisheries. *Env. Res. Lett.*
- Scavia, D., Kelly, E.L.A., Hagy, J.D., 2006. A simple model for forecasting the effects of nitrogen loads on Chesapeake Bay hypoxia. *Estuar. Coasts* 29 (4), 674–684.
- Scavia, D. N.N. Rabalais, R.E. Turner, D. Justic, and W. Wiseman Jr. 2003. Predicting the response of Gulf of Mexico Hypoxia to variations in Mississippi River Nitrogen Load. *Limnol. Oceanogr.* 48(3): 951-956.
- Streeter, H.W. and E.B. Phelps, 1925. A Study in the Pollution and Natural Purification of the Ohio River, III Factors Concerning the Phenomena of Oxidation and Reaeration. US Public Health Service, Public Health Bulletin No. 146, Feb 1925 Reprinted by US PHEW, PHA 1958.
- Sturtz, S., Ligges, U., Gelman, A., 2005. R2winbugs: a package for running WinBUGS from R. *Journal of Statistical Software* 12(3), 1-16.
- Testa, J.M., Clark, J.B., Dennison, W.C., Donovan, E.C., Fisher, A.W., Ni, W., Parker, M., Scavia, D., Spitzer, S.E., Waldrop, A.M., Vargas, V.M.D., Ziegler, G., 2017. Ecological forecasting and the science of hypoxia in Chesapeake Bay. *Bioscience* 67 (7), 614–626.
- US EPA. 2010. Chesapeake Bay total maximum daily load for nitrogen, phosphorus and sediment. Available at: <https://www.epa.gov/chesapeake-bay-tmdl/chesapeake-bay-tmdl-document>.
- WIP 2020. Chesapeake Bay Watershed Implementation Plans. Chesapeake Bay Program. https://www.chesapeakebay.net/what/programs/watershed_implementation. Accessed May 20 2020.