



## Using a Multi-Institutional Ensemble of Watershed Models to Assess Agricultural Conservation Effectiveness in a Future Climate

Haley Kujawa, Margaret Kalcic, Jay Martin, Anna Apostel, Jeffrey Kast, Asmita Murumkar, Grey Evenson, Noel Aloysius , Richard Becker, Chelsie Boles, Remegio Confesor , Awoke Dagneu , Tian Guo, Rebecca Logsdon Muenich , Todd Redder, Yu-Chen Wang, and Donald Scavia

**Research Impact Statement:** Multiple watershed models provide robust predictions of agricultural conservation effectiveness in a future climate.

**ABSTRACT:** This study investigates the combined impacts of climate change and agricultural conservation on the magnitude and uncertainty of nutrient loadings in the Maumee River Watershed, the second-largest watershed of the Laurentian Great Lakes. Two scenarios — baseline agricultural management and increased agricultural conservation — were assessed using an ensemble of five Soil and Water Assessment Tools driven by six climate models. The increased conservation scenario included raising conservation adoption rates from a baseline of existing conservation practices to feasible rates in the near future based on farmer surveys. This increased adoption of winter cover crops on 6%–10% to 60% of cultivated cropland; subsurface placement of phosphorus fertilizers on 35%–60% to 68% of cultivated cropland; and buffer strips intercepting runoff from 29%–34% to 50% of cultivated cropland. Increased conservation resulted in statistically significant ( $p \leq 0.05$ ) reductions in annual loads of total phosphorus (41%), dissolved reactive phosphorus (18%), and total nitrogen (14%) under the highest emission climate scenario (RCP 8.5). While nutrient loads decreased with increased conservation relative to baseline management for all watershed models, different conclusions on the true effectiveness of conservation under climate change may be drawn if only one watershed model was used.

(KEYWORDS: climate change; hydrology; Soil and Water Assessment Tool; nutrients; scenario analysis.)

### INTRODUCTION

An increase in intense phytoplankton blooms since the 1980s has been seen across the globe (Ho et al.

2019), and is expected to worsen with climate change (Paerl and Paul 2012), posing significant risks to human health and ecology (Codd 2000; Codd et al. 2005; Liu et al. 2011; Lee et al. 2017; Wituszynski et al. 2017). A variety of factors contribute to the

Paper No. JAWR-21-0062-P of the *Journal of the American Water Resources Association* (JAWR). Received April 2, 2021; accepted May 19, 2022. © 2022 The Authors. Journal of the American Water Resources Association published by Wiley Periodicals LLC on behalf of American Water Resources Association. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made. **Discussions are open until six months from issue publication.**

Department of Food, Agricultural and Biological Engineering (Kujawa, Kalcic, Martin, Apostel, Kast, Murumkar, Evenson), The Ohio State University Columbus, Ohio, USA; Department of Biomedical, Biological and Chemical Engineering (Aloysius), University of Missouri Columbia, Missouri, USA; Department of Environmental Sciences (Becker), University of Toledo Toledo, Ohio, USA; LimnoTech (Boles; Redder), Ann Arbor, Michigan, USA; Norwegian Institute of Bioeconomy Research (Confesor), Ås, Norway; Environmental Consulting and Technology, Inc. (Dagneu), Ann Arbor, Michigan, USA; Department of Agricultural and Biological Engineering (Guo), Purdue University West Lafayette, Indiana, USA; School of Sustainable Engineering and the Built Environment (Muenich), Arizona State University Tempe, Arizona, USA; and School for Environment and Sustainability (Wang, Scavia), University of Michigan Ann Arbor, Michigan, USA (Correspondence to Kujawa: kujawa.21@osu.edu).

*Citation:* Kujawa, H., M. Kalcic, J. Martin, A. Apostel, Jeffrey Kast, A. Murumkar, G. Evenson, N. Aloysius, R. Becker, C. Boles, R. Confesor, A. Dagneu, T. Guo, Rebecca LogsdonMuenich, T. Redder, Y. Wang, and D. Scavia. 2022. “Using a Multi-Institutional Ensemble of Watershed Models to Assess Agricultural Conservation Effectiveness in a Future Climate.” *JAWRA Journal of the American Water Resources Association* 1–15. <https://doi.org/10.1111/1752-1688.13023>.

intense and often harmful algal blooms (HABs), with a primary driver in many regions being nonpoint source nutrient runoff (Beaver et al. 2014; Stumpf et al. 2016). Scientists often predict climate change will increase nutrient runoff to water bodies with more frequent and more intense precipitation events (Paerl and Paul 2012), but there is evidence that in some regions increased precipitation could be offset by higher evapotranspiration, resulting in decreased nutrient loading with climate change (Kalcic et al. 2019; Kujawa et al. 2020; Scavia et al. 2021).

Watershed models are often used to predict future changes in nutrient loading and assess the influence of conservation practices on water quality (Bosch et al. 2014; Johnson et al. 2015; Verma et al. 2015; Kalcic et al. 2019; Kujawa et al. 2020). However, predictions from watershed models can be highly uncertain due to variability in climate model forecasts (e.g., Kujawa et al. 2020; Miralha et al. 2021) as well as variability in how watershed models simulate load response to landscape changes (e.g., Scavia et al. 2017; Martin et al. 2021). Considerable work in apportioning prediction uncertainty to climate vs. hydrologic models has been done for water quantity in a future climate (Wilby and Harris 2006; Wilby et al. 2006; Kay et al. 2008; Addor et al. 2014). These studies show that climate model uncertainty can greatly outweigh uncertainty from the hydrologic model, or that the hydrologic and climate model uncertainties may be comparable, depending on which factors are considered (e.g., inputs, parameterizations, chosen climate models) (Kay et al. 2008; Bosshard et al. 2013; Karlsson et al. 2016; Thober et al. 2018).

There is no comprehensive body of work on predicting conservation practice effectiveness using an ensemble of climate models, watershed models, and land management scenarios. Of the studies mentioned above, only Karlsson et al. (2016) investigated changes in discharge using multiple climate and watershed models as well as multiple land-use scenarios. The study found land-use change across hydrologic models to significantly affect the extreme hydrologic response (i.e., low flow and flooding), but overall land use had a modest contribution on average discharge variation compared to the climate and watershed models. Kujawa et al. (2020) investigated the uncertainty in predictions of climate change in the Maumee River Watershed using an ensemble of watershed models and found phosphorus predictions could be highly uncertain based on decisions made in the set-up of the watershed model (e.g., subroutines, parameterizations, land management assumptions). While Kujawa et al. (2020) concluded that uncertainty from watershed models is significant for predicting nutrient discharge under a future climate, it may be even more important when the complexity of

land management scenario analysis is added. Scavia et al. (2021) investigated the relative uncertainty of nutrient prediction in a series of climate, watershed, and HABs models, and found the watershed model contributed to the overall uncertainty for nutrient load predictions, albeit less than the HABs and climate models. Few studies have examined the watershed model variations' impact on water quality under changing climate and land management.

The goal of this study was to understand the combined impacts of climate change and increased agricultural conservation (IC) on riverine nutrient loading. The two objectives were (1) to predict if IC will reduce nutrient loadings in a future climate, and (2) to assess whether the effectiveness of agricultural conservation will change between historical and future climate periods. This study was carried out in the Maumee River Watershed, the second-largest watershed of the Great Lakes. We used an ensemble of five watershed models developed using the Soil and Water Assessment Tool (SWAT) and climate data from six downscaled General Circulation Models (GCMs) under the highest emission scenario (RCP 8.5) as well as two agricultural land management scenarios (historical management and IC practices including buffer strips, subsurface placement, and cover crops).

## METHODS

### *Study Area*

The study area was the Maumee River Watershed (~17,300 km<sup>2</sup>), located in northwest Ohio, northeast Indiana, and southeast Michigan (Figure 1). The Maumee River is a major tributary to Lake Erie. Lake Erie has experienced significant eutrophication issues since the 1960s and the region has since focused on managing phosphorus to control eutrophication and HABs (Schindler 1974; De Pinto et al. 1986; Schindler et al. 2016). The 2012 Great Lakes Water Quality Agreements specified phosphorus reduction as the main strategy to control HABs in the western Lake Erie basin (GLWQA 2015; USEPA 2018).

Decreased nutrient loading to Lake Erie is necessary to lessen HABs and protect the public health of the people in the Western Lake Erie Basin. In 2014, Lake Erie bloom toxicity caused the City of Toledo to issue a three-day "Do Not Drink" advisory (Jetoo et al. 2015). While there are few studies directly linking human health and *Microcystis* blooms, some have shown the correlation of increases in liver disease and cancer in areas that have *Microcystis* blooms (Lee et al. 2019; Gorham et al. 2020).




FIGURE 1. Map of the Maumee River Watershed. Soil and Water Assessment Tool (SWAT) models were calibrated to discharge and water quality at the Maumee at Waterville gauge.

The Maumee River Watershed's primary land use is agriculture, specifically row crops such as corn, soy, and winter wheat (Figure 1). The Maumee River watershed is also the second-largest contributor of phosphorus loading to Lake Erie but has the highest phosphorus concentrations which are essential for algal bloom growth (Elser 1999; Michalak et al. 2013). Phosphorus from the Maumee River Watershed is largely from nonpoint sources, primarily agricultural (Maccoux et al. 2016). Therefore, increased attention has been placed on this watershed to reduce phosphorus, particularly from agricultural activities (Scavia et al. 2017; Kalcic et al. 2019; Martin et al. 2021).

### Climate Models

We chose to examine mid-century climate change (2046–2065) because the projections are more certain than for the end-of-century, and agricultural and watershed managers tend to find the early period more relevant for their planning. The timescales of interest, annual and March–July, were chosen because they correspond to loading targets in the Great Lakes Water Quality Agreement as the most relevant predictors of central basin hypoxia and harmful algal bloom size, respectively (GLWQA 2015; Stumpf et al. 2016).

Six GCMs were taken from the Coupled Model Intercomparison Project Phase 5 (CMIP5) ensemble and previously downscaled to  $1/8^\circ$  latitude-longitude ( $\sim 12 \times 12$  km) resolution using Bias-corrected Constructed Analogues (Reclamation 2013). The number of GCMs was chosen based on similar climate and watershed studies (Kay et al. 2008; Velazquez et al. 2013; Prudhomme et al. 2014; Giuntoli et al. 2015; Vetter et al. 2017; Thober et al. 2018). We focused on the highest-emissions scenario, RCP 8.5, and included GCMs that varied across the expected range in precipitation change. Both of these choices would be expected to produce the greatest variation in discharge and nutrient loading at the watershed scale (Michalak et al. 2013; Gao et al. 2019; Kujawa et al. 2020).

The climate model data for the watershed spanned 1.5 standard deviations of the CMIP5 ensemble mean for precipitation and remained close to the mean change (annual mean increase of  $2.76^\circ\text{C}$ ; March–July mean increase of  $2.74^\circ\text{C}$ ) for temperature. The changes in annual precipitation from historical (H; 1996–2015) to mid-century (MC; 2046–2065) ranged from a 5% decrease to a 12% increase, and the mean temperature increased between  $2.5^\circ\text{C}$  and  $3.0^\circ\text{C}$ . The changes in March–July precipitation ranged from a 3% decrease to a 19% increase, and the mean temperature increased between  $2.5^\circ\text{C}$  and  $2.9^\circ\text{C}$  (Table 1).

TABLE 1. CMIP5 climate models and changes from historical (1996–2015) to mid-century (2046–2065).

Shortened climate model name (in text)	Full climate model name	Institute	Reference	Original resolution (°long × °lat)	Annual		March–July	
					Change in temp. (°C)	Change in pre-cip. (%)	Change in temp. (°C)	Change in pre-cip. (%)
CanESM	CanESM2	Canadian Center for Climate Modeling and Analysis	Arora et al. (2011)	2.81° × 2.81°	2.8	12	2.9	12
CSIRO_r6	CSIRO-MK3-6-0	CSIRO Marine and Atmospheric Research	Rotstayn et al. (2010)	1.875° × 1.875°	2.5	5	2.7	19
CSIRO_r4	CSIRO-MK3-4-0	CSIRO Marine and Atmospheric Research	Rotstayn et al. (2010)	1.875° × 1.875°	2.5	2	2.5	4
CSIRO_r10	CSIRO-MK3-10-0	CSIRO Marine and Atmospheric Research	Rotstayn et al. (2010)	1.875° × 1.875°	3.0	13	2.9	17
MPI-ESM	MPI-ESM-LR	Max Planck Institute for Meteorology	Zanchettin et al. (2013)	1.875° × 1.875°	2.5	–5	2.5	–3
NorESM	NorESM1-M	Norwegian Climate Centre	Bentsen et al. (2013)	2.5° × 1.875°	2.9	9	2.7	17

### Watershed Models

The watershed model ensemble was built with SWAT. SWAT is a watershed model commonly used to assess nonpoint source pollution in watersheds, as well as climate change impacts on hydrology and nutrients (e.g., Bosch et al. 2014; Verma et al. 2015; Culbertson et al. 2016; Scavia et al. 2017; Čerkasova et al. 2018; Wang et al. 2018). The model uses inputs of elevation, land use and land cover, climate, and soils, runs on daily time scales, and is able to simulate a wide range of agricultural and land-management practices (Neitsch et al. 2009; Arnold et al. 2012). SWAT has been shown to be a suitable model for the Maumee River Watershed given its ability to represent a range of agricultural management practices and achieve a good calibration (Gebremariam et al. 2014).

Five modeling groups from different institutions built unique SWAT model configurations of the Maumee River Watershed. Some of the inputs, such as point sources and percent of agricultural land with certain management practices, were similar across these models (Table 2). Each SWAT model has the same climate stations and inputs (Figure 1). However, each group independently made most modeling assumptions, such as most land management operations, automatic or manual calibration choices, and specific model routine implementation.

All models were calibrated to a single station (Maumee at Waterville, USGS # 04193500) near the watershed outlet and achieved good standards of performance for discharge and nutrients (Moriassi et al. 2007, 2015; Table S1). However, variations in watershed models that all perform well at the outlet can have significant differences in process representation upstream (Apostel et al. 2021). Hence, we then consider each SWAT model as unique. For more detail on baseline model set-up guidelines and variation among model inputs, see supporting information in Kujawa et al. (2020).

### Agricultural Management Scenarios

The two agricultural management scenarios included baseline management (BM) and IC. Conservation practices used in BM represented historical (2005–2015) rates of cover crops, buffer strips, and subsurface placement/incorporation (Table 2). Cover crops were implemented on 6%–10% of cultivated cropland and buffers intercepted runoff from 29%–34% of cultivated cropland. There was significant variation in subsurface placement, with implementation on 35%–60% of cultivated cropland depending on the model. Interpretation was left to each modeling group on whether to include fertilizer incorporation with tillage as subsurface placement.

TABLE 2. Implementation of agricultural practices considered baseline management (BM) based on guidelines from the literature (see supporting information in Kujawa et al. 2020).

SWAT model (institution where built)	Cover crops	Subsurface placement/incorporation	Buffer strips
UT (University of Toledo)	Cereal rye cover crop planted in 10% of cultivated cropland, limited to corn-soybean rotations	Incorporation via tillage immediately following phosphorus application on 60% of cultivated cropland	Buffers intercepting surface runoff from 32% of cultivated cropland
UM (University of Michigan)	Cereal rye cover crop planted in 8.4% of cultivated cropland	Incorporation via tillage three days after phosphorus application on 57% of cultivated cropland, with 21% of cropland having a mixture of broadcast and incorporation of fertilizer	Buffers intercepting surface runoff from 34% of cultivated cropland
OSU (Ohio State University)	Cereal rye cover crop planted in 8.4% of cultivated cropland and limited to corn-soybean rotations.	Subsurface application of phosphorus fertilizers on 35% of cultivated cropland.	Buffers intercepting surface runoff from 29% of cultivated cropland.
LT (LimnoTech)	Cereal rye cover crop planted in 7.5% of cultivated cropland	Subsurface placement of phosphorus fertilizers on 40% of cultivated cropland	Buffers intercepting surface runoff from 30% of cultivated cropland
HU (Heidelberg University)	Cereal rye cover crop planted in 6% of cultivated cropland	Subsurface placement of phosphorus fertilizers on 43.6% of cultivated cropland	Buffers intercepting surface runoff from 30% of cultivated cropland

The IC scenario increased adoption rates above the BM, from the rates listed in Table 2 to a total rate of cover crops on 60% of cultivated cropland, subsurface placement on 68% of cultivated cropland, and buffer strips intercepting runoff from 50% of cultivated cropland. These increases were chosen in collaboration with a stakeholder advisory group to represent feasible adoptions based on farmer surveys (Martin et al. 2021).

The five SWAT models were run with each management scenario and driven with output from the six climate models, resulting in 60 simulations. We applied downscaled precipitation and temperature outputs to the SWAT models by selecting and directly inputting the climate grid data having the closest centroid to the rain gauge stations included in each SWAT model. All model runs were continuous for 1996–2065 with a 5-year warm-up period beforehand.

#### *Metrics for Assessing Conservation Effectiveness with the Ensemble*

Two objectives were tested through comparisons of these 60 simulations across climate and management scenarios (Figure 2).

Objective 1 assessed changes in nutrient loading due to conservation and climate change. We compared the effect of future climate under the two agricultural management scenarios for each climate and

SWAT model combination. Changes in hydrology and nutrients from average historical (e.g.,  $\overline{BM}_H$ , average of the BM scenario from 1996–2015) to average mid-century (e.g.,  $\overline{IC}_{MC}$ , average of the IC scenario from 2046–2065) climates were calculated for each climate and SWAT model combination as the change under BM in a future climate,

$$\Delta BM_{MC-H} = \frac{\overline{BM}_{MC} - \overline{BM}_H}{\overline{BM}_H} \times 100,$$

and the change due to both IC and future climate,

$$\Delta IC_{MC-H} = \frac{\overline{IC}_{MC} - \overline{BM}_H}{\overline{BM}_H} \times 100.$$

Objective 2 assessed the effectiveness of IC under climate change. We calculated change due to IC in each of the time periods, and the compare them to one another:

$$\Delta IC_H = \frac{\overline{IC}_H - \overline{BM}_H}{\overline{BM}_H} \times 100,$$

$$\Delta IC_{MC} = \frac{\overline{IC}_{MC} - \overline{BM}_{MC}}{\overline{BM}_{MC}} \times 100.$$

These objectives were tested at both annual and March–July timescales.




FIGURE 2. Comparisons of average changes in hydrology and nutrient loads for different land management (BM, IC) and climate (historical [H], mid-century [MC]) scenarios for Objectives 1 and 2.

Results for the change in hydrology under  $\Delta BM_{MC-H}$  were discussed extensively in Kujawa et al. (2020), therefore, this study focuses on the IC scenario. Increased conservation ( $\Delta IC_{MC-H}$ ) was considered effective in a future climate if nutrient loads decreased compared to  $\Delta BM_{MC-H}$ , even if future nutrient loads increased in both  $\Delta IC_{MC-H}$  and  $\Delta BM_{MC-H}$  scenarios. We also assessed if conservation effectiveness changed in a future climate by comparing nutrient loadings for historical climate under BM to historical climate under IC ( $\Delta IC_H$ ) and for mid-century climate under BM to mid-century climate under IC ( $\Delta IC_{MC}$ ) (Figure 2).

The nonparametric Wilcoxon Rank-Sum test was used to test for statistically significant differences between scenarios ( $p \leq 0.05$ ; Wilcoxon 1945). We also used the signal-to-noise ratio, the ensemble mean divided by the interquartile range, to determine whether there was an agreement in direction and magnitude of change within a given scenario. Signal-to-noise greater than one indicates good agreement among models (Giuntoli et al. 2015; Thober et al. 2018).

## RESULTS

### *Changes in Hydrology and Nutrient Loads in a Mid-Century Climate (Objective 1)*

**Annual Changes in Hydrology and Nutrient Loads.** The combination of IC and climate change ( $\Delta IC_{MC-H}$ ) produced no significant differences in annual discharge, subsurface discharge, surface runoff, and evapotranspiration (ET) when compared with the impact of climate change under  $\Delta BM_{MC-H}$  ( $p > 0.05$ ; Figure 3). The mean differences

for the entire ensemble (SWAT and GCMs) in  $\Delta IC_{MC-H}$  were increased discharge (+3%), subsurface discharge (+17%), and ET (+8%), and decreased surface runoff (-15%). The signal-to-noise ratios were similar under both management scenarios (i.e., none changed from below to above one, or vice versa) demonstrating that changes in hydrology were not largely affected by agricultural management choices.

In contrast, differences between the changes in nutrient loads for  $\Delta IC_{MC-H}$  and  $\Delta BM_{MC-H}$  were statistically significant ( $p \leq 0.05$ ). The percent changes in annual loads were less for the  $\Delta IC_{MC-H}$  scenario compared to  $\Delta BM_{MC-H}$ . In  $\Delta IC_{MC-H}$ , total phosphorus (TP) decreased by 41%, dissolved reactive phosphorus (DRP) by 18%, and total nitrogen (TN) by 14%. While DRP and TN signal-to-noise ratios remained below 1, they were greater for  $\Delta IC_{MC-H}$  compared to  $\Delta BM_{MC-H}$ , indicating a tendency toward agreement among ensemble members. The strongest agreement (signal-to-noise >1) was a reduction in TP with climate change under  $\Delta IC_{MC-H}$  (Figure 3).

Some SWAT models predicted increases in annual nutrient loading under  $\Delta IC_{MC-H}$  at mid-century. However, in all cases, there was a lesser increase under  $\Delta IC_{MC-H}$  compared to  $\Delta BM_{MC-H}$  scenario, indicating that IC was always helpful in reducing nutrient loads (Figure 4). While the overall ensemble predicted statistically significant nutrient load reductions between  $\Delta BM_{MC-H}$  and  $\Delta IC_{MC-H}$  (Figure 3), this difference was not always significant for individual SWAT models (Figure 4).

**Changes in March–July Hydrology and Nutrient Loads.** Changes in hydrologic characteristics between  $\Delta BM_{MC-H}$  and  $\Delta IC_{MC-H}$  scenarios for March–July, the period that governs the extent of HABs in Lake Erie’s western basin (GLWQA 2015), were similar to those found at the annual timescale (Figure 5). Climate change alone ( $\Delta BM_{MC-H}$ ) did not result in large changes in discharge (+3%), but had greater changes in surface runoff (-10%), subsurface discharge (+9%), and ET (+12%). The addition of conservation ( $\Delta IC_{MC-H}$ ) resulted in similar and insignificant deviations from  $\Delta BM_{MC-H}$  in discharge (+2%), surface runoff (-11%), subsurface discharge (+7%), and ET (+13%). The only hydrologic change with signal-to-noise above 1 was for ET in both  $\Delta BM_{MC-H}$  and  $\Delta IC_{MC-H}$  (Figure 5).

On the contrary, differences between management scenarios ( $\Delta BM_{MC-H}$  and  $\Delta IC_{MC-H}$ ) were statistically significant ( $p \leq 0.05$ ) for TP, DRP, and TN loadings (Figure 5). DRP increased due to climate change ( $\Delta BM_{MC-H}$  +11%) but decreased with the addition of conservation ( $\Delta IC_{MC-H}$  -11%; Table 3). TP was virtually unchanged due to climate ( $\Delta BM_{MC-H}$ , -2%) and greatly reduced with IC ( $\Delta IC_{MC-H}$ , -34%). Only TP




FIGURE 3. Annual changes for  $\Delta BM_{MC-H}$  and  $\Delta IC_{MC-H}$ . Each boxplot includes the 30 GCM-SWAT combinations. Signal-to-noise is directly above each boxplot. Rank Sum test for statistically significant changes ( $***p \leq 0.001$ ,  $****p \leq 0.0001$ , ns = not significant) between  $\Delta BM_{MC-H}$  and  $\Delta IC_{MC-H}$  denoted above boxplots. DRP, dissolved reactive phosphorus; ET, evapotranspiration, TN, total nitrogen; TP, total phosphorus.

and TN in the  $\Delta IC_{MC-H}$  scenario had signal-to-noise ratios greater than one, signifying ensemble agreement (Figure 5). Taken together, this indicates a greater agreement in a more pronounced effect from IC and climate change than from climate change alone.

Individual SWAT models exhibited March–July load patterns similar to annual loads. All SWAT models had less increase in nutrient loading in the  $\Delta IC_{MC-H}$  scenario compared to the  $\Delta BM_{MC-H}$  scenario under future climate, demonstrating consistency for  $\Delta IC_{MC-H}$  in reducing nutrients. Statistically significant differences were detected between  $\Delta BM_{MC-H}$  and  $\Delta IC_{MC-H}$  for TP in Heidelberg University (HU), LimnoTech (LT), and Ohio State (OSU) SWAT models, for DRP in University of Michigan (UM) and LT SWAT models, and for TN in HU, LT, and OSU SWAT models (Figure 6). Yet, the predicted reductions by the  $\Delta IC_{MC-H}$  scenario for individual SWAT models were not as clear as for the ensemble. Some of the models showed clear improvements under  $\Delta IC_{MC-H}$  relative to  $\Delta BM_{MC-H}$  (e.g., UM), while this change was not as apparent in other models (e.g., UT; Figure 6).

### Changing Effectiveness of Conservation under Climate Change (Objective 2)

The SWAT ensemble showed a slight decrease in the effectiveness of IC for TP and DRP in the mid-century climate (Table 4). However, the differences between mid-century ( $\Delta IC_{MC}$ ) and historical ( $\Delta IC_H$ ) conservation effectiveness were not statistically significant when tested with a two-sided Wilcoxon Rank Sum test ( $p > 0.05$ ; Table S6). On average, IC

reduced annual TP by 40% in historical climate ( $\Delta IC_H$ ), and only 36% by the mid-century ( $\Delta IC_{MC}$ ; Table 4). Similarly, a 24% reduction of DRP in a historical climate dropped to 21% in the mid-century. March–July patterns were similar (Table 4). There was considerable variation among SWAT models in both direction and magnitude of change in conservation effectiveness for phosphorus loading (TP and DRP) by mid-century: HU and LT predicted little to no change ( $\pm 1$ –2 percentage points) in conservation effectiveness in both historical ( $\Delta IC_H$ ) and mid-century ( $\Delta IC_{MC}$ ). OSU and UM showed decreased effectiveness of IC for phosphorus and UT showed increased effectiveness (Table 4). All SWAT models demonstrated greater conservation effectiveness for TN in the mid-century. On average, annual TN was reduced 10% in this historical period ( $\Delta IC_H$ ) and 14% at mid-century ( $\Delta IC_{MC}$ ). Slightly larger TN reductions were produced for March–July, with an 11% reduction in the historical period ( $\Delta IC_H$ ) and 17% in the mid-century ( $\Delta IC_{MC}$ ; Table 4).

## DISCUSSION

### Effects of Climate Change and Increased Conservation on Nutrient Loading

Our goal was to assess the combined impacts of climate change and IC on riverine nutrient loading. The first objective was to use an ensemble of watershed and climate models to assess if IC will reduce nutrient loadings in a future climate ( $\Delta IC_{MC-H}$ ). We




FIGURE 4. Annual results for the individual SWAT models for  $\Delta BM_{MC-H}$  and  $\Delta IC_{MC-H}$ . Rank Sum test for statistically significant changes ( $*p \leq 0.05$ ,  $**p \leq 0.01$ , ns = not significant) between  $\Delta BM_{MC-H}$  and  $\Delta IC_{MC-H}$  denoted above boxplots. Signal to noise values for individual SWAT models can be found in Supporting Information (Table S3).





FIGURE 5. March–July changes for  $\Delta BM_{MC-H}$  and  $\Delta IC_{MC-H}$ . Signal-to-noise ratio is directly above each boxplot. Rank Sum test for statistically significant changes ( $***p \leq 0.001$ ,  $****p \leq 0.0001$ , ns = not significant) between  $\Delta BM_{MC-H}$  and  $\Delta IC_{MC-H}$  denoted above boxplots.

showed that  $\Delta IC_{MC-H}$  would be effective in a future climate, with statistically significant ( $p \leq 0.05$ ) decreases in nutrient loadings relative to  $\Delta BM_{MC-H}$

for both annual and March–July loads of TP (–41% annually; –34%, March–July), DRP (–18% annually; –11% March–July), and TN (–14% annually; –24%

TABLE 3. Average changes for  $\Delta BM_{MC-H}$  and  $\Delta IC_{MC-H}$  between historical (1996–2015) and MC (2046–2065) climate as a percent change  $\pm$  standard deviation (SD).

	TP		DRP		TN	
	Annual	March–July	Annual	March–July	Annual	March–July
$\Delta BM_{MC-H}$	$-7 \pm 13$	$-2 \pm 21$	$+1 \pm 20$	$+11 \pm 26$	$-1 \pm 12$	$-9 \pm 15$
$\Delta IC_{MC-H}$	$-41 \pm 15$	$-34 \pm 21$	$-18 \pm 14$	$-11 \pm 22$	$-14 \pm 15$	$-24 \pm 19$


 FIGURE 6. March–July results for the individual SWAT models for  $\Delta BM_{MC-H}$  and  $\Delta IC_{MC-H}$ . Rank sum test for statistically significant changes ( $*p \leq 0.05$ ,  $**p \leq 0.01$ , ns = not significant) between  $\Delta BM_{MC-H}$  and  $\Delta IC_{MC-H}$  denoted above boxplots. Signal to noise values for individual SWAT models can be found in Supporting Information (Table S3).

March–July; Table 3; Figures 3 and 5). The reductions in annual dissolved and TP loads were larger because there were considerable decreases from December to February (Figure S1). TN, however, exhibited greater percentage decreases during the March–July period because the largest reductions were in March and April (Figure S1).

The potential mid-century reductions in phosphorus and nitrogen due to IC ( $\Delta IC_{MC-H}$  compared to  $\Delta BM_{MC-H}$ ) demonstrate the effectiveness of conservation despite uncertainty associated with climate change. DRP was predicted to have an average, *albeit* small, increase due to climate change alone

( $\Delta BM_{MC-H}$ ); however, introducing IC ( $\Delta IC_{MC-H}$ ) yielded a significant reduction in DRP (Figure 5). Thus, IC should help reduce future HABs because DRP is primarily bioavailable fuel for algal growth (Scavia et al. 2014).

The TP reductions between historical and mid-century were small and highly uncertain under  $\Delta BM_{MC-H}$ , but clear and consistent in  $\Delta IC_{MC-H}$  (Figures 3 and 5). Phosphorus is often prioritized in developing freshwater mitigation strategies because there remains more evidence of phosphorus as the limiting factor in freshwater bloom initiation (Schindler 1974; Schindler et al. 2008, 2016; Stumpf et al. 2016;

TABLE 4. Changes in nutrient loading (%  $\pm$  SD) due to the IC scenario in both historical ( $\Delta IC_H$ ) and mid-century ( $\Delta IC_{MC}$ ) time periods averaged within each SWAT model. For the full list of climate and SWAT models (see Table S3).

Variable	SWAT	Annual		March–July	
		$\Delta IC_H$ (1996–2015)	$\Delta IC_{MC}$ (2046–2065)	$\Delta IC_H$ (1996–2015)	$\Delta IC_{MC}$ (2046–2065)
TP	HU	-42 $\pm$ 1	-39 $\pm$ 2	-37 $\pm$ 2	-36 $\pm$ 1
	LT	-57 $\pm$ 2	-56 $\pm$ 2	-52 $\pm$ 2	-53 $\pm$ 2
	OSU	-35 $\pm$ 2	-25 $\pm$ 3	-29 $\pm$ 2	-21 $\pm$ 3
	UM	-45 $\pm$ 1	-36 $\pm$ 1	-45 $\pm$ 2	-35 $\pm$ 1
	UT	-21 $\pm$ 0	-26 $\pm$ 2	-18 $\pm$ 2	-24 $\pm$ 4
	Average $\pm$ SD	-40 $\pm$ 13	-36 $\pm$ 13	-36 $\pm$ 13	-34 $\pm$ 12
DRP	HU	-19 $\pm$ 1	-18 $\pm$ 1	-17 $\pm$ 1	-16 $\pm$ 1
	LT	-25 $\pm$ 1	-26 $\pm$ 1	-24 $\pm$ 1	-26 $\pm$ 1
	OSU	-24 $\pm$ 2	-17 $\pm$ 1	-16 $\pm$ 1	-11 $\pm$ 1
	UM	-38 $\pm$ 0	-30 $\pm$ 1	-39 $\pm$ 0	-30 $\pm$ 1
	UT	-13 $\pm$ 1	-16 $\pm$ 3	-10 $\pm$ 1	-14 $\pm$ 2
	Average $\pm$ SD	-24 $\pm$ 9	-21 $\pm$ 6	-21 $\pm$ 11	-19 $\pm$ 8
TN	HU	-13 $\pm$ 1	-17 $\pm$ 2	-11 $\pm$ 1	-13 $\pm$ 2
	LT	-17 $\pm$ 1	-26 $\pm$ 2	-19 $\pm$ 2	-30 $\pm$ 4
	OSU	-12 $\pm$ 0	-16 $\pm$ 1	-17 $\pm$ 1	-26 $\pm$ 4
	UM	-7 $\pm$ 1	-7 $\pm$ 1	-6 $\pm$ 1	-8 $\pm$ 1
	UT	-2 $\pm$ 5	-2 $\pm$ 4	-3 $\pm$ 6	-7 $\pm$ 5
	Average $\pm$ SD	-10 $\pm$ 6	-14 $\pm$ 9	-11 $\pm$ 7	-17 $\pm$ 11

USEPA 2018). The predicted reductions of both TP and DRP indicate the conservation practices can decrease nutrient runoff and reduce the extent of Lake Erie HABs now and in the mid-century.

The ensemble predicted a reduction in TN in a mid-century climate under  $\Delta BM_{MC-H}$ , and predicted a further reduction due to  $\Delta IC_{MC-H}$ . Some evidence suggests nitrogen reductions could be critical in limiting freshwater algal bloom size, duration, and toxicity (Chaffin et al. 2014; Gobler et al. 2016; Paerl et al. 2016; Newell et al. 2019). While nitrogen is not at this time considered the top priority nutrient for this watershed (USEPA 2018), it is still important for *Microcystis* bloom duration (Chaffin et al. 2014) and toxicity (Gobler et al. 2016). It is then promising that TN is predicted to be further reduced with IC.

Our second objective was to assess whether the effectiveness of IC will change in a future climate. We found that on average IC was slightly less effective in reducing phosphorus in the mid-century ( $\Delta IC_{MC} < \Delta IC_H$ ), but the difference between these two scenarios was not statistically significant (Table 4; Table S2). Nitrogen was more clearly reduced with conservation in the mid-century ( $\Delta IC_{MC}$ ) because the effects of climate alone reduced TN without additional conservation (Kujawa et al. 2020). Therefore, the combined effects of climate change and IC decreased TN loadings further in the mid-century (Table 4).

Across the two objectives, we found that the combined effects of climate change and IC will likely lead to reductions in nitrogen and phosphorus loading to Lake Erie. Other studies on the climate change in the Maumee River Watershed found reductions in

sediment and nutrients with increased conservation practices. Cousino et al. (2015) simulated 100% no-till on agricultural areas and found this lowered sediment yields by 16% compared to corresponding climate scenarios under historical management with conventional tillage (nutrient data not included in findings). Bosch et al. (2014) evaluated modest adoption rates of agricultural conservation practices: 25% of cropland with cover crops and no-till and 20% with filter strips. They found this scenario of agricultural conservation practices with modest changes in climate showed annual loading reductions of 6% TP, 4% DRP, and 4% TN. Similar to this study, Bosch et al. (2014) also found conservation practices (no-till, cover crops, filter strips) to be less effective in a future climate.

Bosch et al. (2014) and Cousino et al. (2015) chose to focus on variability caused by the different climate scenarios and only include one watershed model. However, our results agree with Bosch et al. (2014) that average nutrient loading in the Maumee can be reduced in a future climate with additional conservation despite increased precipitation, and that a slight decrease in conservation effectiveness may occur with climate change. Furthermore, novel results from this work demonstrated that greater but feasible adoption rates of agricultural conservation practices show greater potential for nutrient reduction. It is important to note while this study demonstrated a need for agricultural conservation to reduce nutrient loadings, some climate change studies in the Maumee found reductions in nutrients under historical management due to climate alone (Kalcic et al. 2019; Scavia et al. 2021).

While most climate scenarios show overall increases in precipitation, it may be important to study the effects of seasonal drought in the Maumee on the impacts of changes in water management. Byun et al. (2019) found decreasing soil moisture in the Great Lakes and Midwest and suggest the combination of drought and temperature stresses may lead to greater irrigation. While the research is lacking, Paul et al. (2020) simulated introducing irrigation to a rain-fed watershed increased surface runoff and suggests it may subsequently increase nutrient loss.

#### *Uncertainty and Variability within the Ensemble of Climate and Watershed Models*

Understanding the extent of uncertainty and variability in watershed modeling and scenario analysis helps to effectively communicate climate change uncertainty to stakeholder groups and inform the scientific development of models (Korfmacher 1998; Gregory and Dieckmann 2013). In this study, there was greater uncertainty in the direction of change in hydrology and water quality when considering climate change alone (under  $\Delta\text{BM}_{\text{MC-H}}$ ) than in the combination of climate change and IC. Depending on the watershed and climate model used, BM results showed increasing or decreasing phosphorus loads in a future climate (Kujawa et al. 2020). However, including increased conservation practices ( $\Delta\text{IC}_{\text{MC-H}}$ ) consistently reduced phosphorus loading as compared to BM ( $\Delta\text{BM}_{\text{MC-H}}$ ). Much of the variability stemmed from differences in setting up and calibrating the models under BM (e.g., management assumptions, spatial discretization of models, and parameters; Evenson et al. 2021; Kujawa et al. 2020) and differences in implementation of the IC scenario (Table 2). Allowing modeling groups to independently develop separate baseline SWAT models allowed for a more holistic accounting of differences in watershed models than if one team was to develop several SWAT models with varying inputs and parameterizations.

While the ensemble demonstrated promising results for greater adoption of conservation practices to reduce nutrients in the future, the confidence in nutrient reduction becomes less apparent if an individual watershed model is chosen. The effectiveness of the  $\Delta\text{IC}_{\text{MC-H}}$  scenario for historical and mid-century climates varied across models, and different conclusions may be reached regarding the impacts of climate change and nutrient reduction potential. For example, the UM model predicts increased phosphorus loadings under  $\Delta\text{BM}_{\text{MC-H}}$  and decreased loadings under  $\Delta\text{IC}_{\text{MC-H}}$ . This creates a clear message that IC is essential to counteract a negative consequence of climate change. However, the UT model showed DRP

reductions under both  $\Delta\text{BM}_{\text{MC-H}}$  and  $\Delta\text{IC}_{\text{MC-H}}$  (Figure 4), suggesting that DRP loadings will decrease with or without additional conservation. While these models send conflicting messages about the value of conservation in future nutrient loading, the ensemble avoids drawing conclusions based on a single watershed model and still captures this variability in watershed model response. We attribute much of the variation of changes in phosphorus loading to the setup and parametrization of the SWAT models (Kujawa et al. 2020). Parameterizations and submodels used in SWAT can affect the dominant transport pathway of phosphorus and the subsequent effects of climate change and agricultural conservation on phosphorus loss.

Communicating model uncertainty and variability to stakeholders is challenging. This study includes many facets of uncertainty in models, such as impacts of parameterization, scenario analysis, and spatial discretization on nutrient predictions in climate analysis. Future research could explicitly investigate each facet of uncertainty and their interactions, as well as other factors, such as emissions scenario, downscaling techniques, watershed model calibration methods (e.g., multi-site calibration), and multiple regions of interest (e.g., Wilby et al. 2006; Kay et al. 2008; Velazquez et al. 2013). Information on the dominant causes of uncertainty in nutrient load prediction can inform subsequent studies on creating a watershed model ensemble that provides a holistic accounting of uncertainty and variability in climate change analysis, as well as inform opportunities for focused watershed model development. Phosphorus management is becoming a critical issue to address concerning eutrophication worldwide (Jeppesen et al. 2009; Bol et al. 2018; Ho et al. 2019), and this study has demonstrated the value of using multiple watershed models to capture uncertainty and variability in scenario results. While the analysis presented herein is limited to one watershed and two land management scenarios, it contributes to a growing body of information on the subject of nutrient modeling and climate analysis, given there will always be some inherent uncertainty regardless of how advanced models become (Beven 2016). One goal is to continuously improve and better integrate climate models, watershed nutrient models, harmful algal boom models, and stakeholder interests to address nutrient runoff and HABs so results can better inform environmental management and adaptation (e.g., Scavia et al. 2021).

Scientists may need guidance to determine what knowledge on model uncertainty is required to further agricultural sustainability in the face of climate change before heading down the “refine-experiment-refine” pathway (Miller et al. 2011). Research has been conducted over the past several decades (e.g.,

Arnell 1999; Moges et al. 2021) in the area of cascading uncertainty in hydrologic variables (e.g., discharge, seven-day low flow, flooding). These studies have contributed valuable information on relative sources of uncertainty (e.g., climate model, hydrologic model, downscaling method) which can guide improved climate impact studies. However, there remains little definitive guidance for watershed management regarding how to deal with the uncertainty of the impacts of climate change on riverine discharge. Therefore, there is still a need to reflect on the practical limits of continuing down the research path of modeling climate uncertainty and agricultural sustainability (Crow 2007). Engaging with diverse stakeholders first to define research questions surrounding climate change uncertainty and watershed modeling could be one way to produce socially robust knowledge that can be used more directly to benefit sustainability in society (Miller et al. 2011).

### *Water Quality Targets and the Future of Lake Erie*

In this study, we found that increasing conservation adoption will likely yield considerable movement toward the load reduction target set for Lake Erie in a mid-century climate, despite uncertainty in climate change and a potential reduction in conservation effectiveness. However, it is critical to note that the phosphorus targets were designed with current climate and lake conditions. Warmer temperatures combined with greater lake stratification could create more ideal situations for HABs to form (Paerl and Paul 2012). For example, Del Giudice et al. (2021) suggested that a 2°C increase in Lake Erie water temperature could lead to blooms that start about 10 days earlier and grow 23% more intense for the same nutrient load. Therefore, the current phosphorus targets may not be sufficient to reduce HABs in a future climate.

The uncertainty in predicting future climate, watershed conditions, and agricultural management, as well as in-lake conditions, complicates management decisions aimed to reduce nutrient loading and HABs. However uncertain these predictions may be, inaction may lead to similar or increased future HABs with negative impacts on Lake Erie's ecology, drinking water, fisheries, and public health (Jetoo et al. 2015; Brooks et al. 2016; Wituszynski et al. 2017).

## CONCLUSIONS

This study used an ensemble of climate and watershed models to predict whether realistic rates of IC

in the Maumee River Watershed, the second largest tributary to Lake Erie, will be effective in reducing nutrients in the mid-century (2046–2065) under the highest emission scenario (RCP 8.5). The IC scenario showed significant reductions in nutrients compared to BM for this period. The combined effects of IC and climate change, on average, predicted annual (March–July) decreases of 41% (34%) for TP, 18% (11%) for DRP, and 14% (24%) for TN. The IC scenario was slightly more effective in reducing phosphorus in the historical period than in the mid-century. In addition, watershed models varied considerably in their assessment of to what degree IC, combined with climate change, will produce phosphorus load reductions. The ensemble consistently predicted nitrogen load reductions due to IC and climate change, in part because climate alone reduced nitrogen loads in the mid-century.

We suggest predictions of hydrology and water quality in a future climate should more frequently employ an ensemble of watershed models. Further study on the effects of individual sources of watershed model uncertainty on predictions of water quality can be used to improve watershed model development and inform future climate impact studies. Interdisciplinary stakeholder engagement should accompany defining further research on model uncertainty and nutrient prediction. In this way, scientists can create a body of literature on model uncertainty more suited to realistically address agricultural sustainability in the face of climate change.

## DATA AVAILABILITY STATEMENT

The data used in this study are available upon reasonable request. Please contact the corresponding author.

## SUPPORTING INFORMATION

Additional supporting information may be found online under the Supporting Information tab for this article: Supplemental data and analysis.

## ACKNOWLEDGMENTS

This project was funded by the Ohio Department of Higher Education Harmful Algal Bloom Research Initiative (R/HAB-5-ODHE). We thank our stakeholder advisory group for their active engagement in designing the project and their review of the work. We also acknowledge the World Climate Research Programme's Working Group on Coupled Modeling, which is responsible for CMIP, the

United States Department of Energy's Program for Climate Model Diagnosis and Intercomparison, and the Global Organization for Earth System Science Portals. We are very appreciative of all research groups involved in the production and distribution of the climate data used in this project (Table 1). Open access funding enabled and organized by ProjektDEAL.

## AUTHOR CONTRIBUTIONS

Haley Kujawa: Data curation; formal analysis; visualization; writing – original draft. Margaret Kalcic: Conceptualization; funding acquisition; methodology; resources; software; supervision; validation; writing – review and editing. Jay Martin: Conceptualization; funding acquisition; methodology; supervision; writing – review and editing. Anna Apostel: Data curation; visualization. Jeffrey Kast: Data curation; writing – review and editing. Asmita Murumkar: Visualization; writing – review and editing. Grey Evenson: Writing – review and editing. Noel Aloysius: Conceptualization; methodology; resources; software; validation. Richard Becker: Software; validation. Chelsie Boles: Software; validation; writing – review and editing. Remegio Confesor: Data curation; formal analysis; writing – review and editing. Awoke Dagne: Software; validation. Tian Guo: Software; validation; writing – review and editing. Rebecca Logsdon Muenich: Software; validation. Todd Redder: Software; validation. Yu-Chen Wang: Resources; software; validation. Donald Scavia: Writing – review and editing.

## LITERATURE CITED

- Addor, N., O. Rössler, N. Köplin, M. Huss, R. Weingartner, and J. Seiber. 2014. "Robust Changes and Sources of Uncertainty in the Projected Hydrological Regimes of Swiss Catchments." *Water Resources Research* 50: 7541–62. <https://doi.org/10.1002/2014WR015549>.
- Apostel, A., M. Kalcic, A. Dagne, G. Evenson, J. Kast, K. King, J. Martin, R.L. Muenich, and D. Scavia. 2021. "Simulating Internal Watershed Processes Using Multiple SWAT Models." *Science of The Total Environment* 759: 143920. <https://doi.org/10.1016/j.scitotenv.2020.143920>.
- Arnell, N.W. 1999. "The Effect of Climate Change on Hydrological Regimes in Europe." *Global Environmental Change* 9: 5–23.
- Arnold, J.G., D.N. Moriasi, P.W. Gassman, K.C. Abbaspour, M.J. White, R. Srinivasan, C. Santhi, et al. 2012. "SWAT: Model Use, Calibration, and Validation." *Transactions of the American Society of Agricultural and Biological Engineers* 55: 1491–508.
- Arora, V.K., J.F. Scinocca, G.J. Boer, J.R. Christian, K.L. Denman, G.M. Flato, V.V. Kharin, W.G. Lee, and W.J. Merryfield. 2011. "Carbon Emission Limits Required to Satisfy Future Representative Concentration Pathways of Greenhouse Gases." *Geophysical Research Letters* 38: 3–8. <https://doi.org/10.1029/2010GL046270>.
- Beaver, J.R., E.E. Manis, K.A. Loftin, J.L. Graham, A.I. Pollard, and R.M. Mitchell. 2014. "Land Use Patterns, Ecoregion, and Microcystin Relationships in U.S. Lakes and Reservoirs: A Preliminary Evaluation." *Harmful Algae* 36: 57–62. <https://doi.org/10.1016/j.hal.2014.03.005>.
- Bentsen, M., I. Bethke, J.B. Debernard, T. Iversen, A. Kirkevåg, Ø. Seland, H. Drange, et al. 2013. "The Norwegian Earth System Model, NorESM1-M — Part 1: Description and Basic Evaluation of the Physical Climate." *Geoscientific Model Development* 6: 687–720. <https://doi.org/10.5194/gmd-6-687-2013>.
- Beven, K. 2016. "Facets of Uncertainty: Epistemic Uncertainty, Non-Stationarity, Likelihood, Hypothesis Testing, and Communication." *Hydrological Sciences Journal* 61: 1652–65. <https://doi.org/10.1080/02626667.2015.1031761>.
- Bol, R., G. Gruau, P.-E. Mellander, R. Dupas, M. Bechmann, E. Skarbøvik, M. Bieroza, et al. 2018. "Challenges of Reducing Phosphorus Based Water Eutrophication in the Agricultural Landscapes of Northwest Europe." *Frontiers in Marine Science* 5: 1–16. <https://doi.org/10.3389/fmars.2018.00276>.
- Bosch, N.S., M.A. Evans, D. Scavia, and J.D. Allan. 2014. "Interacting Effects of Climate Change and Agricultural BMPs on Nutrient Runoff Entering Lake Erie." *Journal of Great Lakes Research* 40: 581–89. <https://doi.org/10.1016/j.jglr.2014.04.011>.
- Bosshard, T., M. Carambia, K. Goergen, S. Kotlarski, P. Krahe, M. Zappa, and C. Schär. 2013. "Quantifying Uncertainty Sources in an Ensemble of Hydrological Climate-Impact Projections." *Water Resources Research* 49: 1523–36. <https://doi.org/10.1029/2011WR011533>.
- Brooks, B.W., J.M. Lazorchak, M.D.A. Howard, M.-V.V. Johnson, S.L. Morton, D.A.K. Perkins, E.D. Reavie, G.I. Scott, S.A. Smith, and J.A. Steevens. 2016. "Are Harmful Algal Blooms Becoming the Greatest Inland Water Quality Threat to Public Health and Aquatic Ecosystems?" *Environmental Toxicology and Chemistry* 35: 6–13. <https://doi.org/10.1002/etc.3220>.
- Byun, K., C.M. Chiu, and A.F. Hamlet. 2019. "Effects of 21st Century Climate Change on Seasonal Flow Regimes and Hydrologic Extremes over the Midwest and Great Lakes Region of the US." *Science of the Total Environment* 650: 1261–77. <https://doi.org/10.1016/j.scitotenv.2018.09.063>.
- Čerkasova, N., G. Umgieser, and A. Ertürk. 2018. "Development of a Hydrology and Water Quality Model for a Large Transboundary River Watershed to Investigate the Impacts of Climate Change — A SWAT Application." *Ecological Engineering* 124: 99–115. <https://doi.org/10.1016/j.ecoleng.2018.09.025>.
- Chaffin, J.D., T.B. Bridgeman, D.L. Bade, and C.N. Moberian. 2014. "Summer Phytoplankton Nutrient Limitation in Maumee Bay of Lake Erie during High- Flow and Low- Flow Years." *Journal of Great Lakes Research* 40: 524–31. <https://doi.org/10.1016/j.jglr.2014.04.009>.
- Codd, G.A. 2000. "Cyanobacterial Toxins, the Perception of Water Quality, and the Prioritisation of Eutrophication Control." *Ecological Engineering* 16: 51–60. [https://doi.org/10.1016/S0925-8574\(00\)00089-6](https://doi.org/10.1016/S0925-8574(00)00089-6).
- Codd, G.A., L.F. Morrison, and J.S. Metcalf. 2005. "Cyanobacterial Toxins: Risk Management for Health Protection." *Toxicology and Applied Pharmacology* 203: 264–72. <https://doi.org/10.1016/j.taap.2004.02.016>.
- Cousino, L., R. Becker, and K. Zmijewski. 2015. "Modeling the Effects of Climate Change on Water, Sediment, and Nutrient Yields from the Maumee River Watershed." *Journal of Hydrology: Regional Studies* 4: 762–75. <https://doi.org/10.1016/j.ejrh.2015.12.039>.
- Crow, M. 2007. "None Dare Call It Hubris: The Limits of Knowledge." *Issues in Science and Technology* 23: 1–4. <https://doi.org/10.1136/jnnp.45.3.284>.
- Culbertson, A.M., J.F. Martin, N. Aloysius, and S.A. Ludsins. 2016. "Anticipated Impacts of Climate Change on 21st Century Maumee River Discharge and Nutrient Loads." *Journal of Great Lakes Research* 42: 1332–42. <https://doi.org/10.1016/j.jglr.2016.08.008>.

- De Pinto, J.V., T.C. Young, and L.M. McIlroy. 1986. "Great Lakes Water Quality Improvement." *Environmental Science & Technology* 20: 752–59. <https://doi.org/10.1021/es00150a001>.
- Del Giudice, D., S. Fang, D. Scavia, T.W. Davis, M.A. Evans, and D.R. Obenour. 2021. "Elucidating Controls on Cyanobacteria Bloom Timing and Intensity via Bayesian Mechanistic Modeling." *Science of the Total Environment* 755: 142487. <https://doi.org/10.1016/j.scitotenv.2020.142487>.
- Elser, J.J. 1999. "The Pathway to Noxious Cyanobacteria Blooms in Lakes: The Food Web as the Final Turn." *Freshwater Biology* 42: 537–43. <https://doi.org/10.1046/j.1365-2427.1999.00471.x>.
- Evenson, G.R., M. Kalcic, Y.C. Wang, D. Robertson, D. Scavia, J. Martin, N. Aloysius, et al. 2021. "Uncertainty in Critical Source Area Predictions from Watershed-Scale Hydrologic Models." *Journal of Environmental Management* 279: 111506. <https://doi.org/10.1016/j.jenvman.2020.111506>.
- Gao, J., A.Y. Sheshukov, H. Yen, K.R. Douglas-Mankin, M.J. White, and J.G. Arnold. 2019. "Uncertainty of Hydrologic Processes Caused by Bias-Corrected CMIP5 Climate Change Projections with Alternative Historical Data Sources." *Journal of Hydrology* 568: 551–61. <https://doi.org/10.1016/j.jhydrol.2018.10.041>.
- Gebremariam, S.Y., J.F. Martin, C. DeMarchi, N.S. Bosch, R. Confesor, and S.A. Ludsin. 2014. "A Comprehensive Approach to Evaluating Watershed Models for Predicting River Flow Regimes Critical to Downstream Ecosystem Services." *Environmental Modelling & Software* 61: 121–34. <https://doi.org/10.1016/j.envsoft.2014.07.004>.
- Giuntoli, I., J.P. Vidal, C. Prudhomme, and D.M. Hannah. 2015. "Future Hydrological Extremes: The Uncertainty from Multiple Global Climate and Global Hydrological Models." *Earth System Dynamics* 6: 267–85. <https://doi.org/10.5194/esd-6-267-2015>.
- GLWQA (Great Lakes Water Quality Agreement). 2015. "Recommended Phosphorus Loading Targets for Lake Erie." Annex 4 Objectives and Targets Task Team Final Report to the Nutrients Annex Subcommittee. <https://www.epa.gov/sites/default/files/2015-06/documents/report-recommended-phosphorus-loading-targets-lake-erie-201505.pdf>.
- Gobler, C.J., J.A.M. Burkholder, T.W. Davis, M.J. Harke, T. Johengen, C.A. Stow, and D.B. Van de Waal. 2016. "The Dual Role of Nitrogen Supply in Controlling the Growth and Toxicity of Cyanobacterial Blooms." *Harmful Algae* 54: 87–97. <https://doi.org/10.1016/j.hal.2016.01.010>.
- Gorham, T., E. Dowling Root, Y. Jia, C.K. Shum, and J. Lee. 2020. "Relationship between Cyanobacterial Bloom Impacted Drinking Water Sources and Hepatocellular Carcinoma Incidence Rates." *Harmful Algae* 95: 101801. <https://doi.org/10.1016/j.hal.2020.101801>.
- Gregory, R., and N. Dieckmann. 2013. "Communicating Uncertainty in Multistakeholder Groups." In *Effective Risk Communication*, edited by Árvai, J., and Rivers, L. III, 56–72. Abingdon, Oxon: Routledge.
- Ho, J.C., A.M. Michalak, and N. Pahlevan. 2019. "Widespread Global Increase in Intense Lake Phytoplankton Blooms Since the 1980s." *Nature* 574: 667–70. <https://doi.org/10.1038/s41586-019-1648-7>.
- Jeppesen, E., B. Kronvang, M. Meerhoff, M. Søndergaard, K.M. Hansen, H.E. Andersen, T.L. Lauridsen, et al. 2009. "Climate Change Effects on Runoff, Catchment Phosphorus Loading and Lake Ecological State, and Potential Adaptations." *Journal of Environmental Quality* 38: 1930–41. <https://doi.org/10.2134/jeq2008.0113>.
- Jetoo, S., V.I. Grover, and G. Krantzberg. 2015. "The Toledo Drinking Water Advisory: Suggested Application of the Water Safety Planning Approach." *Sustain* 7: 9787–808. <https://doi.org/10.3390/su7089787>.
- Johnson, T., J. Butcher, D. Deb, M. Faizullahoy, P. Hummel, J. Kittle, S. Mcginnis, et al. 2015. "Modeling Streamflow and Water Quality Sensitivity to Climate Change and Urban Development in 20 U.S. Watersheds." *Journal of the American Water Resources Association* 51: 1321–41. <https://doi.org/10.1111/1752-1688.12308>.
- Kalcic, M.M., R. Logsdon Muenich, S. Basile, A.L. Steiner, C. Kirchhoff, and D. Scavia. 2019. "Climate Change and Nutrient Loading in the Western Lake Erie Basin: Warming Can Counteract a Wetter Future." *Environmental Science & Technology* 53: 7543–50. <https://doi.org/10.1021/acs.est.9b01274>.
- Karlsson, I.B., T.O. Sonnenborg, J.C. Refsgaard, D. Trolle, C.D. Børgesen, J.E. Olesen, E. Jeppesen, and K.H. Jensen. 2016. "Combined Effects of Climate Models, Hydrological Model Structures and Land Use Scenarios on Hydrological Impacts of Climate Change." *Journal of Hydrology* 535: 301–17. <https://doi.org/10.1016/j.jhydrol.2016.01.069>.
- Kay, A.L., H.N. Davies, V.A. Bell, and R.G. Jones. 2008. "Comparison of Uncertainty Sources for Climate Change Impacts: Flood Frequency in England." *Climatic Change* 92: 41–63. <https://doi.org/10.1007/s10584-008-9471-4>.
- Korfmacher, K.S. 1998. "Water Quality Modeling for Environmental Management: Lessons from the Policy Sciences." *Policy Sciences* 31: 35–54.
- Kujawa, H., M. Kalcic, J. Martin, N. Aloysius, A. Apostel, J. Kast, A. Murumkar, et al. 2020. "The Hydrologic Model as a Source of Nutrient Loading Uncertainty in a Future Climate." *Science of the Total Environment* 724: 138004. <https://doi.org/10.1016/j.scitotenv.2020.138004>.
- Lee, J., S. Lee, and X. Jiang. 2017. "Cyanobacterial Toxins in Freshwater and Food: Important Sources of Exposure to Humans." *Annual Review of Food Science and Technology* 8: 281–304. <https://doi.org/10.1146/annurev-food-030216-030116>.
- Lee, S., J. Kim, B. Choi, G. Kim, and J. Lee. 2019. "Harmful Algal Blooms and Liver Diseases: Focusing on the Areas Near the Four Major Rivers in South Korea." *Journal of Environmental Science and Health, Part C* 37: 356–70. <https://doi.org/10.1080/10590501.2019.1674600>.
- Liu, Y., W. Chen, D. Li, Z. Huang, Y. Shen, and Y. Liu. 2011. "Cyanobacteria/Cyanotoxin-Contaminations and Eutrophication Status Before Wuxi Drinking Water Crisis in Lake Taihu, China." *Journal of Environmental Sciences* 23: 575–81. [https://doi.org/10.1016/S1001-0742\(10\)60450-0](https://doi.org/10.1016/S1001-0742(10)60450-0).
- Maccoux, M.J., A. Dove, S.M. Backus, and D.M. Dolan. 2016. "Total and Soluble Reactive Phosphorus Loadings to Lake Erie." *Journal of Great Lakes Research* 42: 1151–65. <https://doi.org/10.1016/j.jglr.2016.08.005>.
- Martin, J.F., M.M. Kalcic, N. Aloysius, A.M. Apostel, M.R. Brooker, G. Evenson, J.B. Kast, et al. 2021. "Evaluating Management Options to Reduce Lake Erie Algal Blooms Using an Ensemble of Watershed Models." *Journal of Environmental Management* 280: 111710. <https://doi.org/10.1016/j.jenvman.2020.111710>.
- Michalak, A.M., E.J. Anderson, D. Beletsky, S. Boland, N.S. Bosch, T.B. Bridgeman, J.D. Chaffin, et al. 2013. "Record-Setting Algal Bloom in Lake Erie Caused by Agricultural and Meteorological Trends Consistent with Expected Future Conditions." *Proceedings of the National Academy of Sciences of the United States of America* 110: 6448–52. <https://doi.org/10.1073/pnas.1216006110>.
- Miller, T.R., T. Muñoz-Erickson, and C.L. Redman. 2011. "Transforming Knowledge for Sustainability: Towards Adaptive Academic Institutions." *International Journal of Sustainability in Higher Education* 12: 177–92. <https://doi.org/10.1108/1467637111118228>.
- Miralha, L., R.L. Muenich, D. Scavia, K. Wells, A.L. Steiner, M. Kalcic, A. Apostel, S. Basile, and C.J. Kirchhoff. 2021. "Bias Correction of Climate Model Outputs Influences Watershed Model

- Nutrient Load Predictions." *Science of the Total Environment* 759: 143039. <https://doi.org/10.1016/j.scitotenv.2020.143039>.
- Moges, E., Y. Demissie, L. Larsen, and F. Yassin. 2021. "Review: Sources of Hydrological Model Uncertainties and Advances in Their Analysis." *Water* 13: 1–23. <https://doi.org/10.3390/w13010028>.
- Moriassi, D.N., J.G. Arnold, M.W. Van Liew, R.L. Bingner, R.D. Harmel, and T.L. Veith. 2007. "Model Evaluation Guidelines for Systematic Quantification of Accuracy in Watershed Simulations." *American Society of Agricultural and Biological Engineers* 50: 885–900.
- Moriassi, D.N., M.W. Gitau, N. Pai, and P. Daggupati. 2015. "Hydrologic and Water Quality Models: Performance Measures and Evaluation Criteria." *Transactions of the American Society of Agricultural and Biological Engineers* 58: 1763–85. <https://doi.org/10.13031/trans.58.10715>.
- Neitsch, S.L., J.G. Arnold, J.R. Kiniry, and J.R. Williams. 2009. *Soil and Water Assessment Tool Theoretical Documentation Version 2009*. College Station, TX: Soil and Water Research Laboratory.
- Newell, S.E., T.W. Davis, T.H. Johengen, D. Gossiaux, A. Burtner, D. Palladino, and M.J. McCarthy. 2019. "Reduced Forms of Nitrogen are a Driver of Non-Nitrogen-Fixing Harmful Cyanobacterial Blooms and Toxicity in Lake Erie." *Harmful Algae* 81: 86–93. <https://doi.org/10.1016/j.hal.2018.11.003>.
- Paerl, H.W., and V.J. Paul. 2012. "Climate Change: Links to Global Expansion of Harmful Cyanobacteria." *Water Research* 6: 1349–63. <https://doi.org/10.1016/j.watres.2011.08.002>.
- Paerl, H.W., J.T. Scott, M.J. McCarthy, S.E. Newell, W.S. Gardner, K.E. Havens, D.K. Hoffman, S.W. Wilhelm, and W.A. Wurtsbaugh. 2016. "It Takes Two to Tango: When and Where Dual Nutrient (N & P) Reductions Are Needed to Protect Lakes and Downstream Ecosystems." *Environmental Science & Technology* 50: 10805–13. <https://doi.org/10.1021/acs.est.6b02575>.
- Paul, M., S. Dangol, V. Kholodovsky, A.R. Sapkota, M. Negahban-Azar, and S. Lansing. 2020. "Modeling the Impacts of Climate Change on Crop Yield and Irrigation in the Monocacy River Watershed, USA." *Climate* 8: 1–20. <https://doi.org/10.3390/cli8120139>.
- Prudhomme, C., I. Giuntoli, E.L. Robinson, D.B. Clark, N.W. Arnell, R. Dankers, B.M. Fekete, et al. 2014. "Hydrological droughts in the 21st Century, Hotspots and Uncertainties from a Global Multimodel Ensemble Experiment." *Proceedings of the National Academy of Sciences of the United States of America* 111: 3262–67. <https://doi.org/10.1073/pnas.1222473110>.
- Reclamation. 2013. *Downscaled CMIP3 and CMIP5 Climate and Hydrology Projections: Release of Downscaled CMIP5 Climate Projections, Comparison with Preceding Information, and Summary of User Needs*. Denver, CO: U.S. Department of the Interior, Bureau of Reclamation, Technical Services Center.
- Rotstayn, L.D., M.A. Collier, M.R. Dix, Y. Feng, H.B. Gordon, S.P. O'Farrell, I.N. Smith, and J. Syktus. 2010. "Improved Simulation of Australian Climate and ENSO-Related Rainfall Variability in a Global Climate Model with an Interactive Aerosol Treatment." *International Journal of Climatology* 30: 1067–88. <https://doi.org/10.1002/joc.1952>.
- Scavia, D., J. David Allan, K.K. Arend, S. Bartell, D. Beletsky, N.S. Bosch, S.B. Brandt, et al. 2014. "Assessing and Addressing the Re-Eutrophication of Lake Erie: Central Basin Hypoxia." *Journal of Great Lakes Research* 40: 226–46. <https://doi.org/10.1016/j.jglr.2014.02.004>.
- Scavia, D., M. Kalcic, R.L. Muenich, J. Read, N. Aloysius, I. Bertani, C. Boles, et al. 2017. "Multiple Models Guide Strategies for Agricultural Nutrient Reductions." *Frontiers in Ecology and the Environment* 15: 126–32. <https://doi.org/10.1002/fee.1472>.
- Scavia, D., Y.C. Wang, D.R. Obenour, A. Apostel, S.J. Basile, M.M. Kalcic, C.J. Kirchhoff, L. Miralha, R.L. Muenich, and A.L. Steiner. 2021. "Quantifying Uncertainty Cascading from Climate, Watershed, and Lake Models in Harmful Algal Bloom Predictions." *Science of the Total Environment* 759: 143487. <https://doi.org/10.1016/j.scitotenv.2020.143487>.
- Schindler, D.W. 1974. "Eutrophication and Recovery in Experimental Lakes: Implications for Lake Management." *Science* 184: 897–99.
- Schindler, D.W., S.R. Carpenter, S.C. Chapra, R.E. Hecky, and D.M. Orihel. 2016. "Reducing Phosphorus to Curb Lake Eutrophication is a Success." *Environmental Science & Technology* 50: 8923–29. <https://doi.org/10.1021/acs.est.6b02204>.
- Schindler, D.W., R.E. Hecky, D.L. Findlay, M.P. Stainton, B.R. Parker, M.J. Paterson, K.G. Beaty, M. Lyng, and S.E.M. Kasian. 2008. "Eutrophication of Lakes Cannot Be Controlled by Reducing Nitrogen Input: Results of a 37-Year Whole-Ecosystem Experiment." *Proceedings of the National Academy of Sciences of the United States of America* 105: 11254–58. <https://doi.org/10.1073/pnas.0805108105>.
- Stumpf, R.P., L.T. Johnson, T.T. Wynne, and D.B. Baker. 2016. "Forecasting Annual Cyanobacterial Bloom Biomass to Inform Management Decisions in Lake Erie." *Journal of Great Lakes Research* 42: 1174–83. <https://doi.org/10.1016/j.jglr.2016.08.006>.
- Thober, S., R. Kumar, N. Wanders, A. Marx, M. Pan, and O. Rakovec. 2018. "Multi-Model Ensemble Projections of European River Floods and High Flows at 1.5, 2, and 3 Degrees Global Warming." *Environmental Research Letters* 13: 014003. <https://doi.org/10.1088/1748-9326/aa9e35>.
- USEPA. 2018. "U.S. Action Plan for Lake Erie." [https://www.epa.gov/sites/production/files/2018-03/documents/us\\_dap\\_final\\_march\\_1.pdf](https://www.epa.gov/sites/production/files/2018-03/documents/us_dap_final_march_1.pdf).
- Velazquez, J.A., J. Schmid, and S. Ricard. 2013. "An Ensemble Approach to Assess Hydrological Models' Contribution to Uncertainties in the Analysis of Climate Change Impact on Water." *Hydrology and Earth System Sciences* 17: 565–78. <https://doi.org/10.5194/hess-17-565-2013>.
- Verma, S., R. Bhattarai, N.S. Bosch, R.C. Cooke, and P.K. Kalita. 2015. "Climate Change Impacts on Flow, Sediment and Nutrient Export in a Great Lakes Watershed Using SWAT." *Clean Soil Air Water* 43: 1464–74. <https://doi.org/10.1002/clen.201400724>.
- Vetter, T., J. Reinhardt, M. Flörke, A. van Griensven, F. Hattermann, S. Huang, H. Koch, et al. 2017. "Evaluation of Sources of Uncertainty in Projected Hydrological Changes under Climate Change in 12 Large-Scale River Basins." *Climatic Change* 141: 419–33. <https://doi.org/10.1007/s10584-016-1794-y>.
- Wang, Y., J. Bian, Y. Zhao, J. Tang, and Z. Jia. 2018. "Assessment of Future Climate Change Impacts on Nonpoint Source Pollution in Snowmelt Period for a Cold Area Using SWAT." *Scientific Reports* 8: 1–13. <https://doi.org/10.1038/s41598-018-20818-y>.
- Wilby, R.L., and I. Harris. 2006. "A Framework for Assessing Uncertainties in Climate Change Impacts: Low-Flow Scenarios for the River Thames, UK." *Water Resources Research* 42: 1–10. <https://doi.org/10.1029/2005WR004065>.
- Wilby, R.L., P.G. Whitehead, A.J. Wade, D. Butterfield, R.J. Davis, and G. Watts. 2006. "Integrated Modelling of Climate Change Impacts on Water Resources and Quality in a Lowland Catchment: River Kennet, UK." *Journal of hydrology* 330: 204–20. <https://doi.org/10.1016/j.jhydrol.2006.04.033>.
- Wilcoxon, F. 1945. "Individual Comparisons of Grouped Data by Ranking Methods." *Biometrics Bulletin* 1: 80–83. <https://doi.org/10.1093/jee/39.2.269>.
- Wituszynski, D.M., C. Hu, J.F. Martin, J.D. Chaffin, S.A. Ludsin, J. Lee, and F. Zhang. 2017. "Microcystin in Lake Erie Fish: Risk to Human Health and Relationship to Cyanobacterial Blooms." *Journal of Great Lakes Research* 43: 1084–90. <https://doi.org/10.1016/j.jglr.2017.08.006>.
- Zanchettin, D., A. Rubino, D. Matei, O. Bothe, and J.H. Jungclaus. 2013. "Multidecadal-to-Centennial SST Variability in the MPI-ESM Simulation Ensemble for the Last Millennium." *Climate Dynamics* 40: 1301–18. <https://doi.org/10.1007/s00382-012-1361-9>.